

## Product datasheet for RC204112L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TEM8 (ANTXR1) (NM 018153) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** TEM8 (ANTXR1) (NM\_018153) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ATR; GAPO; TEM8 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK NM 018153

**ORF Size:** 999 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC204112).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.


RefSeq: NM 018153.2

RefSeq Size: 2360 bp RefSeq ORF: 1002 bp Locus ID: 84168 **UniProt ID:** Q9H6X2 Cytogenetics: 2p13.3

**Domains: VWA** 

**Protein Families:** Druggable Genome, Transmembrane







**MW:** 37.1 kDa

**Gene Summary:** 

This gene encodes a type I transmembrane protein and is a tumor-specific endothelial marker that has been implicated in colorectal cancer. The encoded protein has been shown to also be a docking protein or receptor for Bacillus anthracis toxin, the causative agent of the disease, anthrax. The binding of the protective antigen (PA) component, of the tripartite anthrax toxin, to this receptor protein mediates delivery of toxin components to the cytosol of cells. Once inside the cell, the other two components of anthrax toxin, edema factor (EF) and lethal factor (LF) disrupt normal cellular processes. Three alternatively spliced variants that encode different protein isoforms have been described. [provided by RefSeq, Oct 2008]