

## Product datasheet for RC204150L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EMAP II (AIMP1) (NM\_004757) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: EMAP II (AIMP1) (NM 004757) Human Tagged ORF Clone Lentiviral Particle

Symbol: EMAP I

**Synonyms:** EMAP2; EMAPII; HLD3; p43; SCYE1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_004757

ORF Size: 936 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC204150).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 004757.2

 RefSeq Size:
 3159 bp

 RefSeq ORF:
 939 bp

 Locus ID:
 9255

 UniProt ID:
 Q12904

 Cytogenetics:
 4q24

Domains: tRNA bind

**Protein Families:** Druggable Genome





MW: 34.4 kDa

**Gene Summary:** 

The protein encoded by this gene is a cytokine that is specifically induced by apoptosis, and it is involved in the control of angiogenesis, inflammation, and wound healing. The release of this cytokine renders the tumor-associated vasculature sensitive to tumor necrosis factor. The precursor protein is identical to the p43 subunit, which is associated with the multi-tRNA synthetase complex, and it modulates aminoacylation activity of tRNA synthetase in normal cells. This protein is also involved in the stimulation of inflammatory responses after proteolytic cleavage in tumor cells. Multiple transcript variants encoding different isoforms have been found for this gene. A pseudogene has been identified on chromosome 20. [provided by RefSeq, Dec 2008]