

Product datasheet for RC204156

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Superoxide Dismutase 3 (SOD3) (NM_003102) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Superoxide Dismutase 3 (SOD3) (NM_003102) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: Superoxide Dismutase 3

Synonyms: EC-SOD

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC204156 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA





Protein Sequence: >RC204156 protein sequence

Red=Cloning site Green=Tags(s)

MLALLCSCLLLAAGASDAWTGEDSAEPNSDSAEWIRDMYAKVTEIWQEVMQRRDDDGTLHAACQVQPSAT LDAAQPRVTGVVLFRQLAPRAKLDAFFALEGFPTEPNSSSRAIHVHQFGDLSQGCESTGPHYNPLAVPHP QHPGDFGNFAVRDGSLWRYRAGLAASLAGPHSIVGRAVVVHAGEDDLGRGGNQASVENGNAGRRLACCVV GVCGPGLWERQAREHSERKKRRRESECKAA


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6213 c04.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_003102

ORF Size: 720 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

ORIGENE

Domains:

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Plasmids are not sterile. For experiments where strict sterility is required, filtration with Note:

0.22um filter is required.

NM 003102.1, NP 003093.1 RefSeq:

RefSeq Size: 1546 bp RefSeq ORF: 723 bp Locus ID: 6649 **UniProt ID:** P08294 4p15.2 Cytogenetics:

sodcu **Protein Families:** Druggable Genome, Secreted Protein

MW: 25.9 kDa

Gene Summary: This gene encodes a member of the superoxide dismutase (SOD) protein family. SODs are


> antioxidant enzymes that catalyze the conversion of superoxide radicals into hydrogen peroxide and oxygen, which may protect the brain, lungs, and other tissues from oxidative stress. Proteolytic processing of the encoded protein results in the formation of two distinct homotetramers that differ in their ability to interact with the extracellular matrix (ECM). Homotetramers consisting of the intact protein, or type C subunit, exhibit high affinity for heparin and are anchored to the ECM. Homotetramers consisting of a proteolytically cleaved form of the protein, or type A subunit, exhibit low affinity for heparin and do not interact with

the ECM. A mutation in this gene may be associated with increased heart disease risk.


[provided by RefSeq, Oct 2015]



Product images:




Circular map for RC204156



Western blot validation of overexpression lysate (Cat# [LY418899]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC204156 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).





Coomassie blue staining of purified SOD3 protein (Cat# [TP304156]). The protein was produced from HEK293T cells transfected with SOD3 cDNA clone (Cat# RC204156) using MegaTran 2.0 (Cat# [TT210002]).