

## Product datasheet for RC204167L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Frizzled 7 (FZD7) (NM\_003507) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Frizzled 7 (FZD7) (NM 003507) Human Tagged ORF Clone Lentiviral Particle

Symbol: FZD7
Synonyms: FzE3

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_003507 **ORF Size:** 1722 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC204167).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 003507.1, NP 003498.1

 RefSeq Size:
 3851 bp

 RefSeq ORF:
 1725 bp

 Locus ID:
 8324

 UniProt ID:
 075084

Cytogenetics: 2q33.1

**Protein Families:** Druggable Genome, Transmembrane





## Frizzled 7 (FZD7) (NM\_003507) Human Tagged ORF Clone Lentiviral Particle - RC204167L4V

**Protein Pathways:** Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling

pathway

MW: 63.4 kDa

**Gene Summary:** Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are

receptors for Wnt signaling proteins. The FZD7 protein contains an N-terminal signal sequence, 10 cysteine residues typical of the cysteine-rich extracellular domain of Fz family members, 7 putative transmembrane domains, and an intracellular C-terminal tail with a PDZ domain-binding motif. FZD7 gene expression may downregulate APC function and enhance beta-catenin-mediated signals in poorly differentiated human esophageal carcinomas.

[provided by RefSeq, Jul 2008]