

Product datasheet for RC204218L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP6V1C2 (NM_144583) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ATP6V1C2 (NM_144583) Human Tagged ORF Clone Lentiviral Particle

Symbol: ATP6V1C2

Synonyms: ATP6C2; VMA5

Mammalian Cell

mmanan Cen

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_144583 **ORF Size:** 1143 bp

ORF Nucleotide

- - 1

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC204218).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 144583.3

RefSeq Size: 3112 bp
RefSeq ORF: 1146 bp
Locus ID: 245973
UniProt ID: Q8NEY4
Cytogenetics: 2p25.1

Domains: V-ATPase C





ATP6V1C2 (NM_144583) Human Tagged ORF Clone Lentiviral Particle - RC204218L4V

Protein Pathways: Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative

phosphorylation, Vibrio cholerae infection

MW: 43.9 kDa

Gene Summary: This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that

mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A,three B, and two G subunits, as well as a C, D, E, F, and H subunit. The V1 domain contains the ATP catalytic site. This gene encodes alternate transcriptional splice variants, encoding different V1 domain C subunit isoforms. [provided by RefSeq, Jul

2008]