

Product datasheet for RC204243L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Tcl1 (TCL1A) (NM_021966) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Tcl1 (TCL1A) (NM_021966) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Tcl1

Synonyms: TCL1

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC204243).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_021966

ORF Size: 342 bp





Tcl1 (TCL1A) (NM_021966) Human Tagged Lenti ORF Clone - RC204243L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 021966.1</u>

 RefSeq Size:
 1410 bp

 RefSeq ORF:
 345 bp

 Locus ID:
 8115

 UniProt ID:
 P56279

Cytogenetics: 14q32.13

Protein Families: Druggable Genome

MW: 13.5 kDa


Gene Summary: Overexpression of the TCL1 gene in humans has been implicated in the development of

mature T cell leukemia, in which chromosomal rearrangements bring the TCL1 gene in close proximity to the T-cell antigen receptor (TCR)-alpha (MIM 186880) or TCR-beta (MIM 186930) regulatory elements (summarized by Virgilio et al., 1998 [PubMed 9520462]). In normal T cells TCL1 is expressed in CD4-/CD8- cells, but not in cells at later stages of differentiation. TCL1 functions as a coactivator of the cell survival kinase AKT (MIM 164730) (Laine et al., 2000


[PubMed 10983986]).[supplied by OMIM, Jul 2010]



Product images:



Circular map for RC204243L1



Double digestion of RC204243L1 using Sgfl and Mlul