

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC204243L2V

## Tcl1 (TCL1A) (NM\_021966) Human Tagged ORF Clone Lentiviral Particle

## Product data:

Product Type:	Lentiviral Particles
Product Name:	Tcl1 (TCL1A) (NM_021966) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Tcl1
Synonyms:	TCL1
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_021966
ORF Size:	342 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204243).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 021966.1</u>
RefSeq Size:	1410 bp
RefSeq ORF:	345 bp
Locus ID:	8115
UniProt ID:	<u>P56279</u>
Cytogenetics:	14q32.13
Protein Families:	Druggable Genome
MW:	13.5 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Overexpression of the TCL1 gene in humans has been implicated in the development of<br/>mature T cell leukemia, in which chromosomal rearrangements bring the TCL1 gene in close<br/>proximity to the T-cell antigen receptor (TCR)-alpha (MIM 186880) or TCR-beta (MIM 186930)<br/>regulatory elements (summarized by Virgilio et al., 1998 [PubMed 9520462]). In normal T cells<br/>TCL1 is expressed in CD4-/CD8- cells, but not in cells at later stages of differentiation. TCL1<br/>functions as a coactivator of the cell survival kinase AKT (MIM 164730) (Laine et al., 2000<br/>[PubMed 10983986]).[supplied by OMIM, Jul 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US