

## Product datasheet for RC204249L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NFX1 (NM\_002504) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** NFX1 (NM\_002504) Human Tagged ORF Clone Lentiviral Particle

Symbol: NFX

Synonyms: NFX2; TEG-42; Tex42

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK ACCN: NM\_002504

ORF Size: 3360 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC204249).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002504.3

 RefSeq Size:
 4721 bp

 RefSeq ORF:
 3363 bp

 Locus ID:
 4799

 UniProt ID:
 Q12986

 Cytogenetics:
 9p13.3

**Domains:** zf-NF-X1, R3H

**Protein Families:** Druggable Genome, Transcription Factors





ORIGENE

**MW:** 124.4 kDa

Gene Summary: MHC class II gene expression is controlled primarily at the transcriptional level by

transcription factors that bind to the X and Y boxes, two highly conserved elements in the proximal promoter of MHC class II genes. The protein encoded by this gene is a transcriptional repressor capable of binding to the conserved X box motif of HLA-DRA and other MHC class II genes in vitro. The protein may play a role in regulating the duration of an inflammatory response by limiting the period in which class II MHC molecules are induced by

IFN-gamma. Three alternative splice variants, each of which encodes a different isoform, have

been identified. [provided by RefSeq, Jul 2008]