

Product datasheet for RC204251L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RPL28 (NM_000991) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RPL28 (NM_000991) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL28
Synonyms: L28

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000991

ORF Size: 411 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204251).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000991.3

 RefSeq Size:
 4245 bp

 RefSeq ORF:
 414 bp

 Locus ID:
 6158

 UniProt ID:
 P46779

 Cytogenetics:
 19q13.42

Domains: Ribosomal_L28e

Protein Pathways: Ribosome





ORIGENE

MW: 15.7 kDa

Gene Summary:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L28E family of ribosomal proteins. It is located in the cytoplasm. Variable expression of this gene in colorectal cancers compared to adjacent normal tissues has been observed, although no correlation between the level of expression and the severity of the disease has been found. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Oct 2008]