

Product datasheet for RC204372L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CALML5 (NM_017422) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CALML5 (NM_017422) Human Tagged ORF Clone Lentiviral Particle

Symbol: CALML5
Synonyms: CLSP

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_017422

ORF Size: 438 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204372).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 017422.3</u>

RefSeq Size: 893 bp
RefSeq ORF: 441 bp
Locus ID: 51806
UniProt ID: Q9NZT1
Cytogenetics: 10p15.1





CALML5 (NM_017422) Human Tagged ORF Clone Lentiviral Particle - RC204372L3V

Protein Pathways: Alzheimer's disease, Calcium signaling pathway, Glioma, GnRH signaling pathway, Insulin

signaling pathway, Long-term potentiation, Melanogenesis, Neurotrophin signaling pathway, Olfactory transduction, Oocyte meiosis, Phosphatidylinositol signaling system, Vascular

smooth muscle contraction

MW: 15.9 kDa

Gene Summary: This gene encodes a novel calcium binding protein expressed in the epidermis and related to

the calmodulin family of calcium binding proteins. Functional studies with recombinant protein demonstrate it does bind calcium and undergoes a conformational change when it does so. Abundant expression is detected only in reconstructed epidermis and is restricted to differentiating keratinocytes. In addition, it can associate with transglutaminase 3, shown to be a key enzyme in the terminal differentiation of keratinocytes. [provided by RefSeq, Jul

2008]