

Product datasheet for RC204404L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SLP76 (LCP2) (NM_005565) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SLP76 (LCP2) (NM_005565) Human Tagged ORF Clone Lentiviral Particle

Symbol: SLP76

Synonyms: IMD81; SLP-76; SLP76

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_005565 **ORF Size:** 1599 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204404).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005565.3</u>

 RefSeq Size:
 2472 bp

 RefSeq ORF:
 1602 bp

 Locus ID:
 3937

 UniProt ID:
 Q13094

 Cytogenetics:
 5q35.1

Domains: SH2, SAM





SLP76 (LCP2) (NM_005565) Human Tagged ORF Clone Lentiviral Particle - RC204404L2V

Protein Pathways: Fc epsilon RI signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor

signaling pathway

MW: 60.2 kDa

Gene Summary: This gene encodes an adapter protein that acts as a substrate of the T cell antigen receptor

(TCR)-activated protein tyrosine kinase pathway. The encoded protein associates with growth factor receptor bound protein 2, and is thought to play a role TCR-mediated intracellular signal transduction. A similar protein in mouse plays a role in normal T-cell development and activation. Mice lacking this gene show subcutaneous and intraperitoneal fetal hemorrhaging,

dysfunctional platelets and impaired viability. [provided by RefSeq, Nov 2016]