

## Product datasheet for **RC204425L2V**

### ATP6V1A (NM\_001690) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	ATP6V1A (NM_001690) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ATP6V1A
Synonyms:	ARCL2D; ATP6A1; ATP6V1A1; DEE93; HO68; IECEE3; VA68; Vma1; VPP2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_001690
ORF Size:	1851 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204425).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001690.2</a>
RefSeq Size:	4607 bp
RefSeq ORF:	1854 bp
Locus ID:	523
UniProt ID:	<a href="#">P38606</a>
Cytogenetics:	3q13.31
Domains:	ATP-synt_ab, ATP-synt_ab_C, ATP-synt_ab_N
Protein Families:	Druggable Genome



[View online »](#)

**Protein Pathways:** Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

**MW:** 68.3 kDa

**Gene Summary:** This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain A subunit isoforms and is found in all tissues. Transcript variants derived from alternative polyadenylation exist. [provided by RefSeq, Jul 2008]