

## Product datasheet for RC204443L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **CLK1 (NM\_004071) Human Tagged ORF Clone Lentiviral Particle**

## **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** CLK1 (NM\_004071) Human Tagged ORF Clone Lentiviral Particle

Symbol: CLK1

Synonyms: CLK; CLK/STY; STY

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_004071 **ORF Size:** 1452 bp

**ORF Nucleotide** 

OTI Disclaimer:

•

Sequence:

The ORF insert of this clone is exactly the same as(RC204443).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 004071.2

 RefSeq Size:
 1933 bp

 RefSeq ORF:
 1455 bp

 Locus ID:
 1195

 UniProt ID:
 P49759

 Cytogenetics:
 2q33.1

**Domains:** pkinase, TyrKc, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase



ORÏGENE

**MW:** 57.3 kDa

**Gene Summary:** 

This gene encodes a member of the CDC2-like (or LAMMER) family of dual specificity protein kinases. In the nucleus, the encoded protein phosphorylates serine/arginine-rich proteins involved in pre-mRNA processing, releasing them into the nucleoplasm. The choice of splice sites during pre-mRNA processing may be regulated by the concentration of transacting factors, including serine/arginine rich proteins. Therefore, the encoded protein may play an indirect role in governing splice site selection. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2009]