

Product datasheet for RC204481L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MEK6 (MAP2K6) (NM_002758) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: MEK6 (MAP2K6) (NM_002758) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: MEK6

Synonyms: MAPKK6; MEK6; MKK6; PRKMK6; SAPKK-3; SAPKK3

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC204481).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_002758

ORF Size: 1002 bp





MEK6 (MAP2K6) (NM_002758) Human Tagged Lenti ORF Clone - RC204481L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 002758.2</u>

 RefSeq Size:
 1879 bp

 RefSeq ORF:
 1005 bp

 Locus ID:
 5608

 UniProt ID:
 P52564

Cytogenetics: 17q24.3

Domains: pkinase, TyrKc, S_TKc

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Fc epsilon RI signaling pathway, GnRH signaling pathway,

MAPK signaling pathway, Toll-like receptor signaling pathway


MW: 37.5 kDa

Gene Summary: This gene encodes a member of the dual specificity protein kinase family, which functions as


a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis. [provided by RefSeq, Jul 2008]



Product images:



Circular map for RC204481L4



Double digestion of RC204481L4 using Sgfl and Mlul