

Product datasheet for RC204485L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Adenosine Receptor A2a (ADORA2A) (NM_000675) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Adenosine Receptor A2a (ADORA2A) (NM_000675) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Adenosine Receptor A2a

Synonyms: A2aR; ADORA2; RDC8

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000675

ORF Size: 1236 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC204485).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000675.3</u>

RefSeq Size: 2583 bp **RefSeq ORF:** 1239 bp

Locus ID: 135

Domains:

UniProt ID: <u>P29274</u>

Cytogenetics: 22q11.23

7tm 1





Adenosine Receptor A2a (ADORA2A) (NM_000675) Human Tagged ORF Clone Lentiviral Particle – RC204485L2V

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Calcium signaling pathway, Neuroactive ligand-receptor interaction, Vascular smooth muscle

contraction

MW: 44.7 kDa

Gene Summary: This gene encodes a member of the guanine nucleotide-binding protein (G protein)-coupled

but it is thought to be non-coding. [provided by RefSeq, Jun 2013]

receptor (GPCR) superfamily, which is subdivided into classes and subtypes. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein, an adenosine receptor of A2A subtype, uses adenosine as the preferred endogenous agonist and preferentially interacts with the G(s) and G(olf) family of G proteins to increase intracellular cAMP levels. It plays an important role in many biological functions, such as cardiac rhythm and circulation, cerebral and renal blood flow, immune function, pain regulation, and sleep. It has been implicated in pathophysiological conditions such as inflammatory diseases and neurodegenerative disorders. Alternative splicing results in multiple transcript variants. A read-through transcript composed of the upstream SPECC1L (sperm antigen with calponin homology and coiled-coil domains 1-like) and ADORA2A (adenosine A2a receptor) gene sequence has been identified,