

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC204523L2V

## WFDC5 (NM\_145652) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	WFDC5 (NM_145652) Human Tagged ORF Clone Lentiviral Particle
Symbol:	WFDC5
Synonyms:	dJ211D12.5; PRG5; WAP1
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_145652
ORF Size:	369 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204523).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 145652.2</u>
RefSeq Size:	1018 bp
RefSeq ORF:	372 bp
Locus ID:	149708
UniProt ID:	Q8TCV5
Cytogenetics:	20q13.12
Protein Families:	Secreted Protein
MW:	13.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a member of the WAP-type four-disulfide core (WFDC) domain family.<br/>Most WFDC proteins contain only one WFDC domain, and this encoded protein contains two<br/>WFDC domains. The WFDC domain, or WAP signature motif, contains eight cysteines forming<br/>four disulfide bonds at the core of the protein, and functions as a protease inhibitor. Most<br/>WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric<br/>and telomeric. This gene belongs to the centromeric cluster. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US