

## Product datasheet for RC204544L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SLC22A17 (NM\_020372) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** SLC22A17 (NM\_020372) Human Tagged ORF Clone Lentiviral Particle

Symbol: SLC22A17

Synonyms: 24p3R; BOCT; BOIT; hBOIT; NGALR; NGALR2; NGALR3

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 020372

ORF Size: 1614 bp

**ORF Nucleotide** 

Judostido The ORE

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC204544).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 020372.2</u>

 RefSeq Size:
 2206 bp

 RefSeq ORF:
 1617 bp

 Locus ID:
 51310

 UniProt ID:
 Q8WUG5

 Cytogenetics:
 14q11.2

**Domains:** sugar\_tr

**Protein Families:** Druggable Genome, Transmembrane





ORIGENE

**MW:** 57.7 kDa

**Gene Summary:** Cell surface receptor for LCN2 (24p3) that plays a key role in iron homeostasis and transport.

Able to bind iron-bound LCN2 (holo-24p3), followed by internalization of holo-24p3 and release of iron, thereby increasing intracellular iron concentration and leading to inhibition of apoptosis. Also binds iron-free LCN2 (apo-24p3), followed by internalization of apo-24p3 and its association with an intracellular siderophore, leading to iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration and resulting in apoptosis (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US