

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC204578L4V

## RNF125 (NM\_017831) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	RNF125 (NM_017831) Human Tagged ORF Clone Lentiviral Particle
Symbol:	RNF125
Synonyms:	TNORS; TRAC-1; TRAC1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_017831
ORF Size:	696 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204578).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 017831.3</u>
RefSeq Size:	5888 bp
RefSeq ORF:	699 bp
Locus ID:	54941
UniProt ID:	<u>Q96EQ8</u>
Cytogenetics:	18q12.1
Domains:	RING
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	RNF125 (NM_017831) Human Tagged ORF Clone Lentiviral Particle – RC204578L4V
Protein Pathways	RIG-I-like receptor signaling pathway
MW:	26.5 kDa
Gene Summary:	This gene encodes a novel E3 ubiquitin ligase that contains a RING finger domain in the N- terminus and three zinc-binding and one ubiquitin-interacting motif in the C-terminus. As a result of myristoylation, this protein associates with membranes and is primarily localized to intracellular membrane systems. The encoded protein may function as a positive regulator in the T-cell receptor signaling pathway. [provided by RefSeq, Mar 2012]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US