

Product datasheet for RC204605L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytochrome P450 2C8 (CYP2C8) (NM 000770) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cytochrome P450 2C8 (CYP2C8) (NM_000770) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cytochrome P450 2C8

CPC8; CYP2C8DM; CYPIIC8; MP-12/MP-20 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 000770 ACCN: **ORF Size:** 1470 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC204605).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000770.3, NP 000761.3

RefSeq Size: 1924 bp RefSeq ORF: 1473 bp Locus ID: 1558 **UniProt ID:** P10632 Cytogenetics: 10q23.33

Domains: p450

Protein Families: Druggable Genome, P450, Transmembrane





Cytochrome P450 2C8 (CYP2C8) (NM_000770) Human Tagged ORF Clone Lentiviral Particle – RC204605L3V

Protein Pathways: Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism,

Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism

MW: 55.8 kDa

Gene Summary: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, benzo(a)pyrene, 7-ethyoxycoumarin, and the anti-cancer drug taxol. This gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Nov

2010]