

## Product datasheet for RC204608L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EDG1 (S1PR1) (NM 001400) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** EDG1 (S1PR1) (NM\_001400) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CD363; CHEDG1; D1S3362; ECGF1; EDG-1; EDG1; S1P1 Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 001400 ACCN: **ORF Size:** 1146 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

**Domains:** 

The ORF insert of this clone is exactly the same as(RC204608).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001400.2

RefSeq Size: 3050 bp RefSeq ORF: 1149 bp Locus ID: 1901 **UniProt ID:** P21453 Cytogenetics: 1p21.2

**Protein Families:** Druggable Genome, GPCR, Transmembrane

7tm 1





## EDG1 (S1PR1) (NM\_001400) Human Tagged ORF Clone Lentiviral Particle - RC204608L1V

**Protein Pathways:** Neuroactive ligand-receptor interaction

MW: 42.8 kDa

**Gene Summary:** The protein encoded by this gene is structurally similar to G protein-coupled receptors and is

highly expressed in endothelial cells. It binds the ligand sphingosine-1-phosphate with high affinity and high specificity, and suggested to be involved in the processes that regulate the differentiation of endothelial cells. Activation of this receptor induces cell-cell adhesion. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016]