

Product datasheet for RC204614L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PYROXD1 (NM_024854) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PYROXD1 (NM_024854) Human Tagged ORF Clone Lentiviral Particle

Symbol: PYROXD²
Synonyms: MFM8

Mammalian Cell F

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_024854

 ORF Size:
 1500 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC204614).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 024854.2

 RefSeq Size:
 4136 bp

 RefSeq ORF:
 1503 bp

 Locus ID:
 79912

 UniProt ID:
 Q8WU10

 Cytogenetics:
 12p12.1

 Domains:
 pyr_redox

Protein Families: Druggable Genome





ORIGENE

MW: 55.8 kDa

Gene Summary:

This gene encodes a nuclear-cytoplasmic pyridine nucleotide-disulphide reductase (PNDR). PNDRs are flavoproteins that catalyze the pyridine nucleotide-dependent reduction of thiol residues in other proteins. The encoded protein belongs to the class I pyridine nucleotide-disulphide oxidoreductase family but lacks the C-terminal dimerization domain found in other family members and instead has a C-terminal nitrile reductase domain. It localizes to the nucleus and to striated sarcomeric compartments. Naturally occurring mutations in this gene cause early-onset myopathy with internalized nuclei and myofibrillar disorganization. A pseudogene of this gene has been defined on chromosome 11. [provided by RefSeq, Apr 2017]