

Product datasheet for RC204709L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hemoglobin subunit gamma 2 (HBG2) (NM_000184) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Hemoglobin subunit gamma 2 (HBG2) (NM_000184) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: Hemoglobin subunit gamma 2

Synonyms: HBG-T1; TNCY

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone

Sequence:

The ORF insert of this clone is exactly the same as(RC204709).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_000184

ORF Size: 441 bp





Hemoglobin subunit gamma 2 (HBG2) (NM_000184) Human Tagged Lenti ORF Clone – RC204709L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000184.2</u>

RefSeq Size: 583 bp RefSeq ORF: 444 bp

Locus ID: 3048

 UniProt ID:
 P69892

 Cytogenetics:
 11p15.4

Domains: globin

MW: 16.1 kDa

Gene Summary: The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen


and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma

product (HBG2) and alanine is found in the A-gamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'- epsilon --

gamma-G -- gamma-A -- delta -- beta--3'. [provided by RefSeq, Jul 2008]



Product images:



Circular map for RC204709L4