

Product datasheet for **RC204745L1V**

UGT (UGT2B4) (NM_021139) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	UGT (UGT2B4) (NM_021139) Human Tagged ORF Clone Lentiviral Particle
Symbol:	UGT2B4
Synonyms:	HLUG25; UDPGT2B4; UDPGTh-1; UDPGTH1; UGT2B11
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_021139
ORF Size:	1584 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204745).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_021139.2
RefSeq Size:	2119 bp
RefSeq ORF:	1587 bp
Locus ID:	7363
UniProt ID:	P06133
Cytogenetics:	4q13.3
Domains:	UDPGT
Protein Families:	Druggable Genome, Transmembrane



[View online »](#)

Protein Pathways: Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

MW: 60.5 kDa

Gene Summary: UDPGTs are of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds. This isozyme is active on polyhydroxylated estrogens (such as estriol, 4-hydroxyestrone and 2-hydroxyestriol) and xenobiotics (such as 4-methylumbelliferone, 1-naphthol, 4-nitrophenol, 2-aminophenol, 4-hydroxybiphenyl and menthol). It is capable of 6 alpha-hydroxyglucuronidation of hyodeoxycholic acid.[UniProtKB/Swiss-Prot Function]