

Product datasheet for RC204745L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UGT (UGT2B4) (NM_021139) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: UGT (UGT2B4) (NM_021139) Human Tagged ORF Clone Lentiviral Particle

Symbol: UGT2B4

Synonyms: HLUG25; UDPGT2B4; UDPGTh-1; UDPGTH1; UGT2B11

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_021139 **ORF Size:** 1584 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204745).

Sequence:
OTI Disclaimer:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 021139.2

 RefSeq Size:
 2119 bp

 RefSeq ORF:
 1587 bp

 Locus ID:
 7363

 UniProt ID:
 P06133

 Cytogenetics:
 4q13.3

Protein Families: Druggable Genome, Transmembrane

UDPGT





UGT (UGT2B4) (NM_021139) Human Tagged ORF Clone Lentiviral Particle - RC204745L2V

Protein Pathways: Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism -

cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and

chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

MW: 60.5 kDa

Gene Summary: UDPGTs are of major importance in the conjugation and subsequent elimination of

potentially toxic xenobiotics and endogenous compounds. This isozyme is active on polyhydroxylated estrogens (such as estriol, 4-hydroxyestrone and 2-hydroxyestriol) and xenobiotics (such as 4-methylumbelliferone, 1-naphthol, 4-nitrophenol, 2-aminophenol, 4-

hydroxybiphenyl and menthol). It is capable of 6 alpha-hydroxyglucuronidation of

hyodeoxycholic acid.[UniProtKB/Swiss-Prot Function]