

## Product datasheet for RC204825L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NMNAT1 (NM\_022787) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** NMNAT1 (NM\_022787) Human Tagged ORF Clone Lentiviral Particle

Symbol: NMNAT1

Synonyms: LCA9; NMNAT; PNAT1; SHILCA

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_022787

ORF Size: 837 bp

**ORF Nucleotide** 

TI. ODE

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC204825).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 022787.2

 RefSeq Size:
 3781 bp

 RefSeq ORF:
 840 bp

 Locus ID:
 64802

 UniProt ID:
 Q9HAN9

 Cytogenetics:
 1p36.22

**Domains:** CTP\_transf\_2

**Protein Pathways:** Metabolic pathways, Nicotinate and nicotinamide metabolism





ORIGENE

**MW:** 31.9 kDa

**Gene Summary:** 

This gene encodes an enzyme which catalyzes a key step in the biosynthesis of nicotinamide adenine dinucleotide (NAD). The encoded enzyme is one of several nicotinamide nucleotide adenylyltransferases, and is specifically localized to the cell nucleus. Activity of this protein leads to the activation of a nuclear deacetylase that functions in the protection of damaged neurons. Mutations in this gene have been associated with Leber congenital amaurosis 9. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are located on chromosomes 1, 3, 4, 14, and 15. [provided by RefSeq, Jul 2014]