

## Product datasheet for **RC204840**

### LILRB1 (NM\_001081639) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	LILRB1 (NM_001081639) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	LILRB1
Synonyms:	CD85J; ILT-2; ILT2; LIR-1; LIR1; MIR-7; MIR7; PIR-B; PIRB
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)



**Protein Sequence:** >RC204840 representing NM\_001081639  
 Red=Cloning site Green=Tags(s)

MTPILTVLICLGLSLGPRTHVQAGHLPKPTLWAEPGSVITQGSPVTLRCQGGQETQEYRLYREKKTAPWI  
 TRIPQELVKKGQFPISITWEHTGRYRCYYGSDTAGRSESSDPLELVVTGAYIKPTLSAQSPVVNSGGN  
 VTLQCDSQVAFDGFILCKEGEDEHPQCLNSQPHARGSSRAIFSVGPVSPSRRWYRCYAYDSNSPYEWSL  
 PSDLLELLVLGVSKKPSLSVQPGPIVAPEETLTLQCGSDAGYNRFVLYKDGEDFLQLAGAQPQAGLSQA  
 NFTLGPVSRSYGQYRCYGAHNLSSEWSAPSDPLDILIAQGFYDRVLSVQPGPTVASGENVTLLCQSQG  
 WMQTFLLTKEGAADDPWRLRSTYQSQKYQAEFPMGPVTSAHAGTYRCYGSQSSKPYLLTHPSDPLELVVS  
 GPSGGPSSPTTGTSTSGPEDQPLTPTGSDPQSGLGRHLGVVIGILVAVILLLLLLLLLLLILRHRHQGK  
 HWTSTQRKADFQHPAGAVGPEPTDRGLQWRSSPAADAQEENLYAAVKHTQPEDGVEMDTRQSPHDEDPA  
 VTVAEVKHSRPREMASPPSPLSGEFLDTKDRQAEEDRQMDTEAAASEAPQDVTYAQLHSLTLRREATEP  
 PPSQEGSPAVPSIYATLAIH


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:**

SgfI-MluI

**Cloning Scheme:**



**Plasmid Map:**


**ACCN:** NM\_001081639

**ORF Size:** 1953 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001081639.3](#), [NP\\_001075108.2](#)

RefSeq Size: 2918 bp

RefSeq ORF: 1956 bp

Locus ID: 10859

UniProt ID: [Q8NHL6](#)

Cytogenetics: 19q13.42

Protein Families: Transmembrane

MW: 71 kDa

**Gene Summary:** This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]