

Product datasheet for RC204849L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD48 (NM_001778) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CD48 (NM_001778) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD48

Synonyms: BCM1; BLAST; BLAST1; hCD48; mCD48; MEM-102; SLAMF2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001778

ORF Size: 729 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204849).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001778.2, NP 001769.2

RefSeq Size: 1155 bp
RefSeq ORF: 732 bp
Locus ID: 962
UniProt ID: P09326

Cytogenetics: 1q23.3

Domains: IG

Protein Families: Druggable Genome, Transmembrane





CD48 (NM_001778) Human Tagged ORF Clone Lentiviral Particle - RC204849L4V

Protein Pathways: Natural killer cell mediated cytotoxicity

MW: 27.7 kDa

Gene Summary: This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which

includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this

gene. [provided by RefSeq, Dec 2011]