

Product datasheet for RC204894L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fas Ligand (FASLG) (NM 000639) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Fas Ligand (FASLG) (NM 000639) Human Tagged ORF Clone Lentiviral Particle

Symbol: Fas Ligand

Synonyms: ALPS1B; APT1LG1; APTL; CD95-L; CD95L; CD178; FASL; TNFSF6; TNLG1A

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000639

ORF Size: 843 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204894).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000639.1

 RefSeq Size:
 1909 bp

 RefSeq ORF:
 846 bp

 Locus ID:
 356

 UniProt ID:
 P48023

Cytogenetics: 1q24.3

Domains: TNF

Protein Families: Druggable Genome, Secreted Protein, Transmembrane





Fas Ligand (FASLG) (NM_000639) Human Tagged ORF Clone Lentiviral Particle - RC204894L4V

Protein Pathways: Allograft rejection, Apoptosis, Autoimmune thyroid disease, Cytokine-cytokine receptor

interaction, Graft-versus-host disease, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Pathways in cancer, Type I diabetes mellitus

MW: 31.5 kDa

Gene Summary: This gene is a member of the tumor necrosis factor superfamily. The primary function of the

encoded transmembrane protein is the induction of apoptosis triggered by binding to FAS. The FAS/FASLG signaling pathway is essential for immune system regulation, including activation-induced cell death (AICD) of T cells and cytotoxic T lymphocyte induced cell death. It has also been implicated in the progression of several cancers. Defects in this gene may be related to some cases of systemic lupus erythematosus (SLE). Alternatively spliced transcript

variants have been described. [provided by RefSeq, Nov 2014]