

Product datasheet for RC204954L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NDUFV1 (NM_007103) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NDUFV1 (NM_007103) Human Tagged ORF Clone Lentiviral Particle

Symbol: NDUFV1

Synonyms: CI-51K; CI51KD; MC1DN4; UQOR1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_007103

 ORF Size:
 1392 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC204954).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 007103.2

 RefSeq Size:
 1631 bp

 RefSeq ORF:
 1395 bp

 Locus ID:
 4723

 UniProt ID:
 P49821

 Cytogenetics:
 11q13.2

Domains: Complex1_51K

Protein Families: Druggable Genome





NDUFV1 (NM_007103) Human Tagged ORF Clone Lentiviral Particle - RC204954L3V

Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease

MW: 50.8 kDa

Gene Summary: The mitochondrial respiratory chain provides energy to cells via oxidative phosphorylation

and consists of four membrane-bound electron-transporting protein complexes (I-IV) and an ATP synthase (complex V). This gene encodes a 51 kDa subunit of the NADH:ubiquinone oxidoreductase complex I; a large complex with at least 45 nuclear and mitochondrial encoded subunits that liberates electrons from NADH and channels them to ubiquinone. This subunit carries the NADH-binding site as well as flavin mononucleotide (FMN)- and Fe-S-biding sites. Defects in complex I are a common cause of mitochondrial dysfunction; a syndrome that occurs in approximately 1 in 10,000 live births. Mitochondrial complex I deficiency is linked to myopathies, encephalomyopathies, and neurodegenerative disorders such as Parkinson's disease and Leigh syndrome. Alternative splicing results in multiple

transcript variants encoding distinct isoforms.[provided by RefSeq, Oct 2009]