

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC204957L4

Cyclin D1 (CCND1) (NM_053056) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cyclin D1 (CCND1) (NM_053056) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Cyclin D1
Synonyms:	BCL1; D11S287E; PRAD1; U21B31
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204957).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I ORF Mlu I GCG ATC GC[C ATG // NNN ACG CGT



ACCN: ORF Size: NM_053056 885 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	yclin D1 (CCND1) (NM_053056) Human Tagged Lenti ORF Clone – RC204957L4
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution M	 chod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 053056.2, NP 444284.1</u>
RefSeq Size:	4304 bp
RefSeq ORF:	888 bp
Locus ID:	595
UniProt ID:	<u>P24385</u>
Cytogenetics:	11q13.3
Domains:	cyclin_C, CYCLIN, cyclin
Protein Families:	Druggable Genome, Stem cell - Pluripotency, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway, Stem cell relevant signaling - Wnt Signaling pathway
Protein Pathway	Acute myeloid leukemia, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Focal adhesion, Glioma, Jak-STAT signaling pathway, Melanoma, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer, Thyroid cancer, Viral myocarditis, Wnt signaling pathway
MW:	33.7 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Science Cyclin D1 (CCND1) (NM_053056) Human Tagged Lenti ORF Clone – RC204957L4

Gene Summary:The protein encoded by this gene belongs to the highly conserved cyclin family, whose
members are characterized by a dramatic periodicity in protein abundance throughout the
cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct
expression and degradation patterns which contribute to the temporal coordination of each
mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4
or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown
to interact with tumor suppressor protein Rb and the expression of this gene is regulated
positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell
cycle progression, are observed frequently in a variety of human cancers. [provided by
RefSeq, Dec 2019]

Product images:



Circular map for RC204957L4

Double digestion of RC204957L4 using Sgfl and Mlul

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US