

Product datasheet for RC204966L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HLA (HLA-DPB1) (NM_002121) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HLA (HLA-DPB1) (NM_002121) Human Tagged ORF Clone Lentiviral Particle

Symbol: HLA-DPB1

Synonyms: DPB1; HLA-DP; HLA-DP1B; HLA-DPB

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 002121

ORF Size: 771 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC204966).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002121.4

 RefSeq Size:
 4055 bp

 RefSeq ORF:
 777 bp

 Locus ID:
 3115

 UniProt ID:
 P04440

 Cytogenetics:
 6p21.32

Domains: MHC_II_beta, ig, IGc1

Protein Families: Transmembrane





HLA (HLA-DPB1) (NM_002121) Human Tagged ORF Clone Lentiviral Particle - RC204966L1V

Protein Pathways: Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid

disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus

erythematosus, Type I diabetes mellitus, Viral myocarditis

MW: 29.2 kDa

Gene Summary: HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a

heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different

molecules. [provided by RefSeq, Jul 2008]