

## Product datasheet for RC205065L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RAG2 (NM\_000536) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type: Lentiviral Particles

Product Name: RAG2 (NM 000536) Human Tagged ORF Clone Lentiviral Particle

Symbol: RAG2 Synonyms: RAG-2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_000536

 ORF Size:
 1581 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC205065).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000536.1, NP 000527.1

 RefSeq Size:
 2457 bp

 RefSeq ORF:
 1584 bp

 Locus ID:
 5897

 UniProt ID:
 P55895

**Protein Families:** Druggable Genome

**Protein Pathways:** Primary immunodeficiency

11p12







**MW:** 59.2 kDa

**Gene Summary:** 

This gene encodes a protein that is involved in the initiation of V(D)J recombination during B and T cell development. This protein forms a complex with the product of the adjacent recombination activating gene 1, and this complex can form double-strand breaks by cleaving DNA at conserved recombination signal sequences. The recombination activating gene 1 component is thought to contain most of the catalytic activity, while the N-terminal of the recombination activating gene 2 component is thought to form a six-bladed propeller in the active core that serves as a binding scaffold for the tight association of the complex with DNA. A C-terminal plant homeodomain finger-like motif in this protein is necessary for interactions with chromatin components, specifically with histone H3 that is trimethylated at lysine 4. Mutations in this gene cause Omenn syndrome, a form of severe combined immunodeficiency associated with autoimmune-like symptoms. [provided by RefSeq, Jul 2008]