

## Product datasheet for RC205074L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Lactate Dehydrogenase B (LDHB) (NM\_002300) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Lactate Dehydrogenase B (LDHB) (NM\_002300) Human Tagged ORF Clone Lentiviral Particle

Symbol: Lactate Dehydrogenase B

Synonyms: HEL-S-281; LDH-B; LDH-H; LDHBD; TRG-5

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_002300

**ORF Size:** 1002 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC205074).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002300.3

 RefSeq Size:
 1336 bp

 RefSeq ORF:
 1005 bp

 Locus ID:
 3945

 UniProt ID:
 P07195

 Cytogenetics:
 12p12.1

Domains: ldh

**Protein Families:** Druggable Genome





## Lactate Dehydrogenase B (LDHB) (NM\_002300) Human Tagged ORF Clone Lentiviral Particle – RC205074L1V

**Protein Pathways:** Cysteine and methionine metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways,

Propanoate metabolism, Pyruvate metabolism

MW: 36.5 kDa

**Gene Summary:** This gene encodes the B subunit of lactate dehydrogenase enzyme, which catalyzes the

interconversion of pyruvate and lactate with concomitant interconversion of NADH and NAD+ in a post-glycolysis process. Alternatively spliced transcript variants have been found for this gene. Recent studies have shown that a C-terminally extended isoform is produced by use of

an alternative in-frame translation termination codon via a stop codon readthrough

mechanism, and that this isoform is localized in the peroxisomes. Mutations in this gene are associated with lactate dehydrogenase B deficiency. Pseudogenes have been identified on

chromosomes X, 5 and 13. [provided by RefSeq, Feb 2016]