

Product datasheet for **RC205092**

RACK1 (NM_006098) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: RACK1 (NM_006098) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: RACK1
Synonyms: Gnb2-rs1; GNB2L1; H12.3; HLC-7; PIG21
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >RC205092 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGACTGAGCAGATGACCCTTCGTGGCACCTCAAGGGCCACAACGGCTGGTAACCCAGATCGCTACTA
CCCCGCAGTCCCGGACATGATCCTCTCCGCCTCTCGAGATAAGACCATCATCATGTGGAACTGACCAG
GGATGAGACCAACTATGGAATCCACAGCGTGCTCTGCGGGTCACTCCCACTTTGTTAGTGATGTGGTT
ATCTCCTCAGATGGCCAGTTTGCCTCTCAGGCTCCTGGGATGGAACCTGCGCCTCTGGGATCTCAAA
CGGGCACCAACGAGGCGATTTGTGGCCATACCAAGGATGTGCTGAGTGTGGCCTTCTCTCTGACAA
CCGGCAGATTGTCTCTGGATCTCGAGATAAAACCATCAAGCTATGGAATACCCCTGGGTGTGTGCAAATAC
ACTGTCCAGGATGAGAGCCACTCAGAGTGGGTGTCTTGTGTCCGCTTCTCGCCCAACAGCAGCAACCCTA
TCATCGTCTCCTGTGGCTGGGACAAGCTGGTCAAGGATGGAACCTGGCTAACTGCAAGCTGAAGACCAA
CCACATTGGCCACACAGGCTATCTGAACACGGTGACTGTCTCTCCAGATGGATCCCTCTGTGCTTCTGGA
GGCAAGGATGGCCAGGCCATGTTATGGGATCTCAACGAAGGCAACACCTTTACACGCTAGATGGTGGGG
ACATCATCAACGCCCTGTGCTTCAGCCCTAACCGCTACTGGCTGTGTGCTGCCACAGGCCCCAGCATCAA
GATCTGGGATTTAGAGGGAAGATCATTGTAGATGAAGCAAGCAAGTTATCAGTACCAGCAGCAAG
GCAGAACCACCCAGTGACCTCCCTGGCCTGGTCTGCTGATGGCCAGACTCTGTTTGTGGCTACACGG
ACAACCTGGTGGGAGTGTGGCAGGTGACCATTGGCACACGC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC205092 protein sequence
 Red=Cloning site Green=Tags(s)

MTEQMTLRGTLKGHNGWVTQIATTPQFPDMILSASRDKTIIMWKLTRDETNYGIPQRALRGHSHFVSDVV
 ISSDGFALSGSWDGLRLWDLTTGTTTRRFVGHTKDVLVSAFSSDNRQIVSGSRDKTIKLWNTLGVCKY
 TVQDESHSEWVSCVRFSPNSSNP IIVSCGWDKLKVVWNLANCKLKTNHIGHTGYLNTVTVSPDGSLCASG
 GKDGQAMLWDLNEGKHL YTLDDGGDIINALCFSPNRYWLC AATGPSIKIWDLEGKIIIVDELKQEVISTSSK
 AEPPQCTSLAWSADGQTLFAGYTDNLVVRVWQVTIGTR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Chromatograms: https://cdn.origene.com/chromatograms/mk6061_e01.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:




ACCN: NM_006098


| | |
|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ORF Size: | 951 bp |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM_006098.5 |
| RefSeq Size: | 1125 bp |
| RefSeq ORF: | 954 bp |
| Locus ID: | 10399 |
| UniProt ID: | P63244 |
| Cytogenetics: | 5q35.3 |
| Domains: | WD40 |
| Protein Families: | Druggable Genome |
| MW: | 35.1 kDa |

Gene Summary:


Scaffolding protein involved in the recruitment, assembly and/or regulation of a variety of signaling molecules. Interacts with a wide variety of proteins and plays a role in many cellular processes. Component of the 40S ribosomal subunit involved in translational repression (PubMed:23636399). Involved in the initiation of the ribosome quality control (RQC), a pathway that takes place when a ribosome has stalled during translation, by promoting ubiquitination of a subset of 40S ribosomal subunits (PubMed:28132843). Binds to and stabilizes activated protein kinase C (PKC), increasing PKC-mediated phosphorylation. May recruit activated PKC to the ribosome, leading to phosphorylation of EIF6. Inhibits the activity of SRC kinases including SRC, LCK and YES1. Inhibits cell growth by prolonging the G0/G1 phase of the cell cycle. Enhances phosphorylation of BMAL1 by PRKCA and inhibits transcriptional activity of the BMAL1-CLOCK heterodimer. Facilitates ligand-independent nuclear translocation of AR following PKC activation, represses AR transactivation activity and is required for phosphorylation of AR by SRC. Modulates IGF1R-dependent integrin signaling and promotes cell spreading and contact with the extracellular matrix. Involved in PKC-dependent translocation of ADAM12 to the cell membrane. Promotes the ubiquitination and proteasome-mediated degradation of proteins such as CLEC1B and HIF1A. Required for VANGL2 membrane localization, inhibits Wnt signaling, and regulates cellular polarization and oriented cell division during gastrulation. Required for PTK2/FAK1 phosphorylation and dephosphorylation. Regulates internalization of the muscarinic receptor CHRM2. Promotes apoptosis by increasing oligomerization of BAX and disrupting the interaction of BAX with the anti-apoptotic factor BCL2L. Inhibits TRPM6 channel activity. Regulates cell surface expression of some GPCRs such as TBXA2R. Plays a role in regulation of FLT1-mediated cell migration. Involved in the transport of ABCB4 from the Golgi to the apical bile canalicular membrane (PubMed:19674157). Promotes migration of breast carcinoma cells by binding to and activating RHOA (PubMed:20499158).[UniProtKB/Swiss-Prot Function]

Product images:

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY GNB2L1 (Cat# RC205092, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GNB2L1 (Cat# [TA502717]). Positive lysates [LY401838] (100ug) and [LC401838] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY401838]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC205092 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified RACK1 protein (Cat# [TP305092]). The protein was produced from HEK293T cells transfected with RACK1 cDNA clone (Cat# RC205092) using MegaTran 2.0 (Cat# [TT210002]).