

Product datasheet for RC205130L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cbl c (CBLC) (NM_012116) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cbl c (CBLC) (NM_012116) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cbl of

Synonyms: CBL-3; CBL-SL; RNF57

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_012116

 ORF Size:
 1422 bp

ORF Nucleotide

1 122 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC205130).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 012116.2

 RefSeq Size:
 1591 bp

 RefSeq ORF:
 1425 bp

 Locus ID:
 23624

 UniProt ID:
 Q9ULV8

 Cytogenetics:
 19q13.32

Protein Families: Druggable Genome





Cbl c (CBLC) (NM_012116) Human Tagged ORF Clone Lentiviral Particle - RC205130L1V

Protein Pathways: Chronic myeloid leukemia, Endocytosis, ErbB signaling pathway, Insulin signaling pathway,

Jak-STAT signaling pathway, Pathways in cancer, T cell receptor signaling pathway, Ubiquitin

mediated proteolysis

MW: 52.5 kDa

Gene Summary: This gene encodes a member of the Cbl family of E3 ubiquitin ligases. Cbl proteins play

important roles in cell signaling through the ubiquitination and subsequent downregulation

of tyrosine kinases. Expression of this gene may be restricted to epithelial cells, and

alternatively spliced transcript variants encoding multiple isoforms have been observed for

this gene. [provided by RefSeq, Feb 2012]