

Product datasheet for RC205243L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PIP5K2 alpha (PIP4K2A) (NM 005028) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PIP5K2 alpha (PIP4K2A) (NM 005028) Human Tagged ORF Clone Lentiviral Particle

Symbol: PIP5K2 alpha

Synonyms: PI5P4KA; PIP5K2A; PIP5KII-alpha; PIP5KIIA; PIPK

Mammalian Cell

. . .

Puromycin

Selection: Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_005028 **ORF Size:** 1218 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC205243).

Sequence:

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

The molecular sequence of this clone aligns with the gene accession number as a point of

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005028.3

RefSeq Size: 3833 bp
RefSeq ORF: 1221 bp
Locus ID: 5305

UniProt ID: P48426

Cytogenetics: 10p12.2

Domains: PIP5K

Protein Families: Druggable Genome





Protein Pathways: Inositol phosphate metabolism, Phosphatidylinositol signaling system, Regulation of actin

cytoskeleton

MW: 46.2 kDa

ORÏGENE

Gene Summary: Phosphatidylinositol-5,4-bisphosphate, the precursor to second messengers of the

phosphoinositide signal transduction pathways, is thought to be involved in the regulation of secretion, cell proliferation, differentiation, and motility. The protein encoded by this gene is one of a family of enzymes capable of catalyzing the phosphorylation of phosphatidylinositol-5-phosphate on the fourth hydroxyl of the myo-inositol ring to form phosphatidylinositol-5,4-bisphosphate. The amino acid sequence of this enzyme does not show homology to other kinases, but the recombinant protein does exhibit kinase activity. This gene is a member of the phosphatidylinositol-5-phosphate 4-kinase family. [provided by RefSeq, Jul 2008]