

Product datasheet for RC205243L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PIP5K2 alpha (PIP4K2A) (NM_005028) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: PIP5K2 alpha (PIP4K2A) (NM 005028) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: PIP5K2 alpha

Synonyms: PI5P4KA; PIP5K2A; PIP5KII-alpha; PIP5KIIA; PIPK

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide


The ORF insert of this clone is exactly the same as(RC205243).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_005028

ORF Size: 1218 bp





PIP5K2 alpha (PIP4K2A) (NM_005028) Human Tagged Lenti ORF Clone - RC205243L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 005028.3</u>

 RefSeq Size:
 3833 bp

 RefSeq ORF:
 1221 bp

 Locus ID:
 5305

 UniProt ID:
 P48426

Cytogenetics: 10p12.2

Domains: PIP5K

Protein Families: Druggable Genome

Protein Pathways: Inositol phosphate metabolism, Phosphatidylinositol signaling system, Regulation of actin

cytoskeleton


MW: 46.2 kDa

Gene Summary: Phosphatidylinositol-5,4-bisphosphate, the precursor to second messengers of the

phosphoinositide signal transduction pathways, is thought to be involved in the regulation of secretion, cell proliferation, differentiation, and motility. The protein encoded by this gene is one of a family of enzymes capable of catalyzing the phosphorylation of phosphatidylinositol-5-phosphate on the fourth hydroxyl of the myo-inositol ring to form phosphatidylinositol-5,4-bisphosphate. The amino acid sequence of this enzyme does not show homology to other kinases, but the recombinant protein does exhibit kinase activity. This gene is a member of the phosphatidylinositol-5-phosphate 4-kinase family. [provided by RefSeq, Jul 2008]



Product images:



Circular map for RC205243L4