

Product datasheet for RC205288L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCSF Receptor (CSF1R) (NM_005211) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MCSF Receptor (CSF1R) (NM 005211) Human Tagged ORF Clone Lentiviral Particle

Symbol: MCSF Receptor

Synonyms: BANDDOS; C-FMS; CD115; CSF-1R; CSFR; FIM2; FMS; HDLS; M-CSF-R

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_005211 **ORF Size:** 2916 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205288).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005211.2, NP 005202.2

 RefSeq Size:
 3985 bp

 RefSeq ORF:
 2919 bp

 Locus ID:
 1436

 UniProt ID:
 P07333

Cytogenetics: 5q32

Domains: pkinase, TyrKc, S_TKc, ig, IGc2, IG

Protein Families: Druggable Genome, Protein Kinase, Transmembrane





Protein Pathways: Cytokine-cytokine receptor interaction, Endocytosis, Hematopoietic cell lineage, Pathways in

cancer

MW: 107.8 kDa

Gene Summary: The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine

which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL), HL cell lines

and anaplastic large cell lymphoma. [provided by RefSeq, Mar 2017]