

Product datasheet for RC205320L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Monoamine Oxidase B (MAOB) (NM_000898) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Monoamine Oxidase B (MAOB) (NM_000898) Human Tagged ORF Clone Lentiviral Particle

Symbol: MAOB **Mammalian Cell**

Selection:

None

1560 bp

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: ACCN: NM_000898 **ORF Size:**

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205320).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

> reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000898.3

RefSeq Size: 2611 bp RefSeq ORF: 1563 bp Locus ID: 4129 **UniProt ID:** P27338 **Cytogenetics:** Xp11.3

Domains: Amino_oxidase

Protein Families: Druggable Genome, Transmembrane





Monoamine Oxidase B (MAOB) (NM_000898) Human Tagged ORF Clone Lentiviral Particle – RC205320L1V

Protein Pathways: Arginine and proline metabolism, Drug metabolism - cytochrome P450, Glycine, serine and

threonine metabolism, Histidine metabolism, Metabolic pathways, Phenylalanine metabolism,

Tryptophan metabolism, Tyrosine metabolism

MW: 58.8 kDa

Gene Summary: The protein encoded by this gene belongs to the flavin monoamine oxidase family. It is a

enzyme located in the mitochondrial outer membrane. It catalyzes the oxidative deamination

of biogenic and xenobiotic amines and plays an important role in the metabolism of neuroactive and vasoactive amines in the central nervous sysytem and peripheral tissues. This protein preferentially degrades benzylamine and phenylethylamine. [provided by

RefSeq, Jul 2008]