

## Product datasheet for RC205322L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GIRK1 (KCNJ3) (NM\_002239) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: GIRK1 (KCNJ3) (NM\_002239) Human Tagged ORF Clone Lentiviral Particle

Symbol: GIRK1

Synonyms: GIRK1; KGA; KIR3.1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM\_002239

ORF Size: 1503 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC205322).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002239.2</u>

 RefSeq Size:
 4740 bp

 RefSeq ORF:
 1506 bp

 Locus ID:
 3760

 UniProt ID:
 P48549

 Cytogenetics:
 2q24.1

Domains: IRK

**Protein Families:** Druggable Genome, Ion Channels: Potassium, Transmembrane





**MW:** 56.7 kDa

**Gene Summary:** 

Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. The encoded protein, which has a greater tendency to allow potassium to flow into a cell rather than out of a cell, is controlled by G-proteins and plays an important role in regulating heartbeat. It associates with three other G-protein-activated potassium channels to form a heteromultimeric pore-forming complex that also couples to neurotransmitter receptors in the brain and whereby channel activation can inhibit action potential firing by hyperpolarizing the plasma membrane. These multimeric G-protein-gated inwardly-rectifying potassium (GIRK) channels may play a role in the pathophysiology of epilepsy, addiction, Down's syndrome, ataxia, and Parkinson's disease. Alternative splicing results in multiple transcript variants encoding distinct proteins. [provided by RefSeq, May 2012]