

Product datasheet for RC205324L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ID2 (NM_002166) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ID2 (NM_002166) Human Tagged ORF Clone Lentiviral Particle

Symbol: ID2

Synonyms: bHLHb26; GIG8; ID2A; ID2H

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002166

ORF Size: 402 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205324).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002166.4</u>

 RefSeq Size:
 1402 bp

 RefSeq ORF:
 405 bp

 Locus ID:
 3398

 UniProt ID:
 Q02363

 Cytogenetics:
 2p25.1

Domains: HLH





ID2 (NM_002166) Human Tagged ORF Clone Lentiviral Particle - RC205324L4V

Protein Families: ES Cell Differentiation/IPS, Stem cell relevant signaling - TGFb/BMP signaling pathway,

Transcription Factors

Protein Pathways: TGF-beta signaling pathway

MW: 14.9 kDa

Gene Summary: The protein encoded by this gene belongs to the inhibitor of DNA binding family, members of

which are transcriptional regulators that contain a helix-loop-helix (HLH) domain but not a basic domain. Members of the inhibitor of DNA binding family inhibit the functions of basic helix-loop-helix transcription factors in a dominant-negative manner by suppressing their heterodimerization partners through the HLH domains. This protein may play a role in negatively regulating cell differentiation. A pseudogene of this gene is located on

chromosome 3. [provided by RefSeq, Aug 2011]