

Product datasheet for RC205338

MAGEB2 (NM 002364) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: MAGEB2 (NM_002364) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: MAGEB2

Synonyms: CT3.2; DAM6; MAGE-XP-2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC205338 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC205338 protein sequence

Red=Cloning site Green=Tags(s)

MPRGQKSKLRAREKRRKARDETRGLNVPQVTEAEEEEAPCCSSSVSGGAASSSPAAGIPQKPQRAPTTAA AAAAGVSSTKSKKGAKSHQGEKNASSSQASTSTKSPSEDPLTRKSGSLVQFLLYKYKIKKSVTKGEMLKI VGKRFREHFPEILKKASEGLSVVFGLELNKVNPNGHTYTFIDKVDLTDEESLLSSWDFPRRKLLMPLLGV IFLNGNSATEEEIWEFLNMLGVYDGEEHSVFGEPWKLITKDLVQEKYLEYKQVPSSDPPRFQFLWGPRAY AETSKMKVLEFLAKVNGTTPCAFPTHYEEALKDEEKAGV


TRTRPLEOKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6316 a09.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_002364

ORF Size: 957 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>



MAGEB2 (NM_002364) Human Tagged ORF Clone - RC205338

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

35.3 kDa

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 002364.5</u>

 RefSeq Size:
 1628 bp

 RefSeq ORF:
 960 bp

 Locus ID:
 4113

 UniProt ID:
 015479

Cytogenetics: Xp21.2

MW:

Gene Summary: This gene is a member of the MAGEB gene family. The members of this family have their


entire coding sequences located in the last exon, and the encoded proteins show 50 to 68% sequence identity to each other. The promoters and first exons of the MAGEB genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. This gene is localized in the DSS (dosage-sensitive sex reversal) critical region. It is expressed in testis and placenta, and

in a significant fraction of tumors of various histological types. The MAGEB genes are


clustered on chromosome Xp22-p21. [provided by RefSeq, Jul 2008]



Product images:





Circular map for RC205338



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MAGEB2 (Cat# RC205338, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAGEB2 (Cat# [TA811621])(1:2000). Positive lysates [LY419377] (100ug) and [LC419377] (20ug) can be purchased separately from OriGene.







Western blot validation of overexpression lysate (Cat# [LY419377]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC205338 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified MAGEB2 protein (Cat# [TP305338]). The protein was produced from HEK293T cells transfected with MAGEB2 cDNA clone (Cat# RC205338) using MegaTran 2.0 (Cat# [TT210002]).