

Product datasheet for **RC205341L3V**

BAF53A (ACTL6A) (NM_178042) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	BAF53A (ACTL6A) (NM_178042) Human Tagged ORF Clone Lentiviral Particle
Symbol:	BAF53A
Synonyms:	ACTL6; Arp4; ARPN-BETA; BAF53A; INO80K
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_178042
ORF Size:	1290 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC205341).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_178042.2
RefSeq Size:	1956 bp
RefSeq ORF:	1164 bp
Locus ID:	86
UniProt ID:	O96019
Cytogenetics:	3q26.33
Protein Families:	Druggable Genome, Transcription Factors
MW:	47.5 kDa



[View online »](#)

Gene Summary:

This gene encodes a family member of actin-related proteins (ARPs), which share significant amino acid sequence identity to conventional actins. Both actins and ARPs have an actin fold, which is an ATP-binding cleft, as a common feature. The ARPs are involved in diverse cellular processes, including vesicular transport, spindle orientation, nuclear migration and chromatin remodeling. This gene encodes a 53 kDa subunit protein of the BAF (BRG1/brm-associated factor) complex in mammals, which is functionally related to SWI/SNF complex in *S. cerevisiae* and *Drosophila*; the latter is thought to facilitate transcriptional activation of specific genes by antagonizing chromatin-mediated transcriptional repression. Together with beta-actin, it is required for maximal ATPase activity of BRG1, and for the association of the BAF complex with chromatin/matrix. Three transcript variants that encode two different protein isoforms have been described. [provided by RefSeq, Jul 2008]