

## Product datasheet for RC205380L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rb2 p130 (RBL2) (NM\_005611) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Rb2 p130 (RBL2) (NM\_005611) Human Tagged ORF Clone Lentiviral Particle

 Symbol:
 Rb2 p130

 Synonyms:
 P130; Rb2

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM\_005611

ORF Size: 3417 bp

**ORF Nucleotide** 

3117 bp

Sequence:

Cytogenetics:

The ORF insert of this clone is exactly the same as(RC205380).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005611.2

 RefSeq Size:
 4903 bp

 RefSeq ORF:
 3420 bp

 Locus ID:
 5934

 UniProt ID:
 Q08999

Domains: RB\_B, RB\_A, CYCLIN

**Protein Families:** Druggable Genome, Transcription Factors

16q12.2





**Protein Pathways:** Cell cycle, TGF-beta signaling pathway

MW: 128.3 kDa

**Gene Summary:** Key regulator of entry into cell division. Directly involved in heterochromatin formation by

maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone

methyltransferases KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Probably acts as a transcription repressor by recruiting chromatin-modifying enzymes to promoters. Potent inhibitor of E2F-mediated trans-activation, associates preferentially with E2F5. Binds to cyclins A and E. Binds to and may be involved in the transforming capacity of the adenovirus E1A protein. May act as a

tumor suppressor.[UniProtKB/Swiss-Prot Function]