

## Product datasheet for RC205382L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Chromogranin C (SCG2) (NM 003469) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Chromogranin C (SCG2) (NM\_003469) Human Tagged ORF Clone Lentiviral Particle

Symbol: Chromogranin C

CHGC; EM66; SgII; SN Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 003469 ACCN:

**ORF Size:** 1851 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC205382).

OTI Disclaimer:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 003469.3

RefSeq Size: 2586 bp RefSeq ORF: 1854 bp Locus ID: 7857 **UniProt ID:** P13521 Cytogenetics: 2q36.1

Granin **Protein Families:** Secreted Protein, Transmembrane





## Chromogranin C (SCG2) (NM\_003469) Human Tagged ORF Clone Lentiviral Particle – RC205382L3V

**MW:** 70.7 kDa

**Gene Summary:** 

The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. Studies in rodents suggest that the full-length protein, secretogranin II, is involved in the packaging or sorting of peptide hormones and neuropeptides into secretory vesicles. The full-length protein is cleaved to produce the active peptide secretoneurin, which exerts chemotaxic effects on specific cell types, and EM66, whose function is unknown. [provided by RefSeq, Jul 2008]