

## Product datasheet for RC205431L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## UCMA (NM\_145314) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** UCMA (NM\_145314) Human Tagged ORF Clone Lentiviral Particle

Symbol: UCMA

**Synonyms:** C10orf49; GRP; GRP/UCMA

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 145314

ORF Size: 414 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC205431).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 145314.1, NP 660357.2

RefSeq Size: 840 bp
RefSeq ORF: 417 bp
Locus ID: 221044
UniProt ID: Q8WVF2
Cytogenetics: 10p13

**Protein Families:** Secreted Protein

**MW:** 16.6 kDa







## **Gene Summary:**

This gene encodes a chondrocyte-specific, highly charged protein that is abundantly expressed in the upper immature zone of fetal and juvenile epiphyseal cartilage. The encoded protein undergoes proteolytic processing to generate a mature protein that is secreted into the extracellular matrix. The glutamic acid residues in the encoded protein undergo gamma carboxylation in a vitamin K-dependent manner. Undercarboxylation of the encoded protein is associated with osteoarthritis in humans. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2015]