

Product datasheet for RC205434L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Activin A Receptor Type IC (ACVR1C) (NM 145259) Human Tagged ORF Clone Lentiviral **Particle**

Product data:

Product Type: Lentiviral Particles

Product Name: Activin A Receptor Type IC (ACVR1C) (NM_145259) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Activin A Receptor Type IC

ACVRLK7; ALK7 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag:

ACCN: NM 145259

ORF Size: 1479 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC205434).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 145259.1, NP 660302.1

RefSeq Size: 3267 bp RefSeq ORF: 1482 bp Locus ID: 130399 **UniProt ID:** Q8NER5 Cytogenetics: 2q24.1

Protein Families: Druggable Genome, Protein Kinase, Transmembrane





Activin A Receptor Type IC (ACVR1C) (NM_145259) Human Tagged ORF Clone Lentiviral Particle – RC205434L3V

Protein Pathways: Adherens junction, Chronic myeloid leukemia, Colorectal cancer, Endocytosis, MAPK signaling

pathway, Pancreatic cancer, Pathways in cancer, TGF-beta signaling pathway

MW: 54.7 kDa

Gene Summary: ACVR1C is a type I receptor for the TGFB (see MIM 190180) family of signaling molecules.

Upon ligand binding, type I receptors phosphorylate cytoplasmic SMAD transcription factors, which then translocate to the nucleus and interact directly with DNA or in complex with other transcription factors (Bondestam et al., 2001 [PubMed 12063393]).[supplied by OMIM, Mar

2008]