

Product datasheet for **RC205473L2V**

RASA1 (NM_002890) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	RASA1 (NM_002890) Human Tagged ORF Clone Lentiviral Particle
Symbol:	RASA1
Synonyms:	CM-AVM; CMAVM; CMAVM1; GAP; p120; p120GAP; p120RASGAP; PKWS; RASA; RASGAP
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_002890
ORF Size:	3141 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC205473).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_002890.1
RefSeq Size:	4402 bp
RefSeq ORF:	3144 bp
Locus ID:	5921
UniProt ID:	P20936
Cytogenetics:	5q14.3
Domains:	C2, SH2, SH3, PH, RasGAP
Protein Families:	Druggable Genome



[View online »](#)

Protein Pathways: Axon guidance, MAPK signaling pathway

MW: 116.4 kDa

Gene Summary: The protein encoded by this gene is located in the cytoplasm and is part of the GAP1 family of GTPase-activating proteins. The gene product stimulates the GTPase activity of normal RAS p21 but not its oncogenic counterpart. Acting as a suppressor of RAS function, the protein enhances the weak intrinsic GTPase activity of RAS proteins resulting in the inactive GDP-bound form of RAS, thereby allowing control of cellular proliferation and differentiation. Mutations leading to changes in the binding sites of either protein are associated with basal cell carcinomas. Mutations also have been associated with hereditary capillary malformations (CM) with or without arteriovenous malformations (AVM) and Parkes Weber syndrome. Alternative splicing results in two isoforms where the shorter isoform, lacking the N-terminal hydrophobic region but retaining the same activity, appears to be abundantly expressed in placental but not adult tissues. [provided by RefSeq, May 2012]