

Product datasheet for RC205480L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RPL37A (NM_000998) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RPL37A (NM_000998) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL37/ Synonyms: L37A

Mammalian Cell None

Selection:

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_000998

ORF Size: 276 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205480).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000998.4

RefSeq Size: 434 bp
RefSeq ORF: 279 bp
Locus ID: 6168
UniProt ID: P61513
Cytogenetics: 2q35

Domains: Ribosomal_L37ae

Protein Pathways: Ribosome





RPL37A (NM_000998) Human Tagged ORF Clone Lentiviral Particle - RC205480L1V

MW: 10.3 kDa

Gene Summary:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L37AE family of ribosomal proteins. It is located in the cytoplasm. The protein contains a C4-type zinc finger-like domain. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]