

## Product datasheet for RC205494L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Non Neuronal Enolase (ENO1) (NM 001428) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Non Neuronal Enolase (ENO1) (NM\_001428) Human Tagged ORF Clone Lentiviral Particle

Symbol: Non Neuronal Enolase

Synonyms: ENO1L1; HEL-S-17; MPB1; NNE; PPH

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_001428 **ORF Size:** 1302 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC205494).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001428.2

RefSeq Size: 2204 bp
RefSeq ORF: 1305 bp
Locus ID: 2023
UniProt ID: P06733
Cytogenetics: 1p36.23
Domains: enolase

**Protein Families:** Druggable Genome, Transcription Factors





## Non Neuronal Enolase (ENO1) (NM\_001428) Human Tagged ORF Clone Lentiviral Particle – RC205494L2V

**Protein Pathways:** Glycolysis / Gluconeogenesis, Metabolic pathways, RNA degradation

**MW:** 47.2 kDa

**Gene Summary:** This gene encodes alpha-enolase, one of three enolase isoenzymes found in mammals. Each

isoenzyme is a homodimer composed of 2 alpha, 2 gamma, or 2 beta subunits, and functions as a glycolytic enzyme. Alpha-enolase in addition, functions as a structural lens protein (taucrystallin) in the monomeric form. Alternative splicing of this gene results in a shorter isoform that has been shown to bind to the c-myc promoter and function as a tumor suppressor. Several pseudogenes have been identified, including one on the long arm of chromosome 1. Alpha-enolase has also been identified as an autoantigen in Hashimoto encephalopathy.

[provided by RefSeq, Jan 2011]